Author Index

AHMED A.E.E., VEITCH J. & WHALEY K. Mechanism of action of an inhibitor of complement-mediated prevention of immune precipitation, 139
AHMED A.E.E. *see* SANDILANDS G.P.
ALBECK M. *see* SREDNI B.
ALBEROLA-ILA J. *see* LOZANO F.
ALBERTI S. *see* TOYE P.G.
ALEKSANDERSEN M., HEIN W.R., LANDSVERK T. & MCCLURE S. Distribution of lymphocyte subsets in the large intestinal lymphoid follicles of lambs, 391
ALLEN J.I. *see* KAY N.E.
ALLINGTON M.P. *see* DUTIA B.M.
AMLOT P.L. *see* DEVEY M.E.
ANGELUCCI J. *see* MALAN BOREL I.
ANNEMA A. *see* SLUITER W.
ANTONI F. *see* MASON D.
APPLETON J.A. *see* CARLISLE M.S.
ARDAILLOU N. *see* SAVIC V.
ARDAILLOU R. *see* SAVIC V.
ARENA A. *see* HORMAECHE C.E.
ASKONAS B.A. *see* PEMBERTON R.M.
ATKINSON J.P. *see* PURCELL D.F.J.
ATKINSON Y.H., MURRAY A.W., KRILIS S., VADAS M.A. & LOPEZ A.F. Human tumour necrosis factor-alpha (TNF-α) directly stimulates arachidonic acid release in human neutrophils, 82
AUBERGER P., SONTHONNAX S., PEYRON J.-F., MARI B. & FEHLMANN M. A chymotryptic-type serine protease is required for IL-2 production by Jurkat T cells, 547
AURBACH D. *see* SREDNI B.
AYALA A., PERRIN M.M. & CHAUDRY I.H. Defective macrophage antigen presentation following haemorrhage is associated with the loss of MHC class II (Ia) antigens, 33

BABIUK L.A. *see* FITZPATRICK D.R.
BARAT A. *see* GONZÁLEZ-CABRERO J.
BARNETT B.C., HARTLMAYR I., GRAHAM C.M. & THOMAS D.B. Single amino acid residues in a synthetic peptide of influenza haemagglutinin, HA 1 177-199, distinguish I-A^d- and I-E^d-restricted T-cell epitopes, 48
BASSENGE E. *see* HAUSCHILDT S.
BATE C.A.W., TAVERNE J., DAVÉ A. & PLAYFAIR J.H.L. Malaria exoantigens induce T-independent antibody that blocks their ability to induce TNF, 315
BEILEFELDT OHMANN H. *see* FITZPATRICK D.R.
BENSAID A.M. *see* TOYE P.G.
BESKORWAYNE T. *see* FITZPATRICK D.R.
BESSLER W. *see* HAUSCHILDT S.
BINAGHI R.A. *see* MALAN BOREL I.
BIRD P. *see* CALVERT J.E.
BIRD P. *see* DEVEY M.E.
BISCHOFF P. *see* MOOG C.
BLAAS P., *see* YAMAMOTO H.
BLEASDALE-BARR K.M. *see* DEVEY M.E.
BLOOM E.T. *see* UMEHARA H.
BORCA M.V. *see* SRIKUMARAN S.
BOYD R.L. *see* GODFREY D.I.
BOYD R.L. *see* WILSON T.J.
BRACH M. *see* REIDEL D.
BRONDZ B.D. *see* ZAITSEVA M.B.
BROUWER M. *see* DE JONG R.
BUJDOSO R. *see* DUTIA B.M.
BUSSE R. *see* HAUSCHILDT S.
BUTTKE T.M. *see* TEBBEY P.W.

CAHILL R.N.P. *see* WITHERDEN D.A.
CALVERT J.E., JOHNSTONE R., DUGGAN-KEEN M.F. & BIRD P. Immunoglobulin G subclasses secreted by human B cells *in vitro* in response to interleukin-2 and polyclonal activators, 162
CAMPBELL P.A. *see* STEDMAN K.E.
CARLISLE M.S., MCGREGOR D.D. & APPLETON J.A. The role of mucus in antibody-mediated rapid expulsion of *Trichinella spiralis* in suckling rats, 126
CHAO T.-Y., OHNISHI H. & CHU T.M. Indirect inhibition of generation of murine lymphokine-activated killer cell activity in splenocyte cultures by interferon-gamma, 116
CHAPMAN C.J. *see* JORDAN R.K.
CHAUDRY I.H. *see* AYALA A.
CHEERS C., ZHAN Y.F. & EGAN P.J. *In vivo* IL-1 potentiates both specific and non-specific arms of immune response to infection, 411
CHEERS C. *see* EGAN P.
CHILD J.A. *see* JONES R.A.
CHOW S.C. & JONDAL M. A central role for phosphoinositide hydrolysis in activating the lytic mechanisms of human natural killer cells, 106
CHU T.M. *see* CHAO T.-Y.
COMPSTON D.A.S. *see* SCOLDING N.J.
CROATTO M. *see* PHILLIPS W.A.

DAHA M.R. *see* SPECHT GRIJP-GLANDORF R.J.A.M.
DAVÉ A. *see* BATE C.A.W.
DAVRINCHE C. *see* DEMARES M.J.
DAY M.J. *see* MACPHEE I.A.M.
DEACON N.J. *see* PURCELL D.F.J.
DE HEER E. *see* SPECHT GRIJP-GLANDORF R.J.A.M.
DE JONG R., BROUWER M., REBEL V.I., VAN SEVENTER G.A., MIEDEMA F. & VAN LIER R.A.W. Generation of alloreactive cytolytic T lymphocytes by immobilized anti-CD3 monoclonal antibodies. Analysis of requirements for human cytolytic T-lymphocyte differentiation, 357
DEKKER-NOOREN C.C.E.M. *see* SPECHT GRIJP-GLANDORF R.J.A.M.
DEMARES M.J. & DAVRINCHE C. Prevalence of the 1·8 kb complement factor H mRNA in human lung, 150
DEVEY M.E., BLEASDALE-BARR K.M., BIRD P. & AMLOT P.L. Antibodies of different human IgG subclasses show distinct patterns of affinity maturation after immunization with keyhole limpet haemocyanin, 168
DING L. *see* ZHANG Y.H.
DOE W.F. *see* PAVLI P.
DOUGAN G. *see* TITE J.P.
DUGGAN-KEEN M.F. *see* CALVERT J.E.
DUTTA B.M., HOPKINS J., ALLINGTON M.P., BUJDOSO R. & MCCONNELL I. Characterization of monoclonal antibodies specific for α- and β-chains of sheep MHC class II, 27

EARLY E. *see* LAURENCE J.
EGAN P. & CHEERS C. Relationship between colony-stimulating activity and interferon production during infection, 191
EGAN P.J. *see* CHEERS C.
EGIDO J. *see* GONZÁLEZ-CABRERO J.
ELZENGA-CLAASEN I. *see* SLUITER W.

FEHLMANN M. *see* AUBERGER P.
FERNANDEZ-RUIZ E. *see* SOMOZA C.
FERRY B.L., STARKEY P.M., SARGENT I.L., WATT G.M.O., JACKSON M. & REDMAN C.W.G. Cell populations in the human early pregnancy decidua: natural killer activity and response to interleukin-2 of CD56-positive large granular lymphocytes, 446
FINBLOOM D.S. *see* PURI R.K.

Author Index

AHMED A.E.E., VEITCH J. & WHALEY K. Mechanism of action of an inhibitor of complement-mediated prevention of immune precipitation, 139
AHMED A.E.E. *see* SANDILANDS G.P.
ALBECK M. *see* SREDNI B.
ALBEROLA-ILA J. *see* LOZANO F.
ALBERTI S. *see* TOYE P.G.
ALEKSANDERSEN M., HEIN W.R., LANDSVERK T. & MCCLURE S. Distribution of lymphocyte subsets in the large intestinal lymphoid follicles of lambs, 391
ALLEN J.I. *see* KAY N.E.
ALLINGTON M.P. *see* DUTIA B.M.
AMLOT P.L. *see* DEVEY M.E.
ANGELUCCI J. *see* MALAN BOREL I.
ANNEMA A. *see* SLUITER W.
ANTONI F. *see* MASON D.
APPLETON J.A. *see* CARLISLE M.S.
ARDAILLOU N. *see* SAVIC V.
ARDAILLOU R. *see* SAVIC V.
ARENA A. *see* HORMAECHE C.E.
ASKONAS B.A. *see* PEMBERTON R.M.
ATKINSON J.P. *see* PURCELL D.F.J.
ATKINSON Y.H., MURRAY A.W., KRILIS S., VADAS M.A. & LOPEZ A.F. Human tumour necrosis factor-alpha (TNF-α) directly stimulates arachidonic acid release in human neutrophils, 82
AUBERGER P., SONTHONNAX S., PEYRON J.-F., MARI B. & FEHLMANN M. A chymotryptic-type serine protease is required for IL-2 production by Jurkat T cells, 547
AURBACH D. *see* SREDNI B.
AYALA A., PERRIN M.M. & CHAUDRY I.H. Defective macrophage antigen presentation following haemorrhage is associated with the loss of MHC class II (Ia) antigens, 33

BABIUK L.A. *see* FITZPATRICK D.R.
BARAT A. *see* GONZÁLEZ-CABRERO J.
BARNETT B.C., HARTLMAYR I., GRAHAM C.M. & THOMAS D.B. Single amino acid residues in a synthetic peptide of influenza haemagglutinin, HA 1 177-199, distinguish I-A^d- and I-E^d-restricted T-cell epitopes, 48
BASSENGE E. *see* HAUSCHILDT S.
BATE C.A.W., TAVERNE J., DAVÉ A. & PLAYFAIR J.H.L. Malaria exoantigens induce T-independent antibody that blocks their ability to induce TNF, 315
BEILEFELDT OHMANN H. *see* FITZPATRICK D.R.
BENSAID A.M. *see* TOYE P.G.
BESKORWAYNE T. *see* FITZPATRICK D.R.
BESSLER W. *see* HAUSCHILDT S.
BINAGHI R.A. *see* MALAN BOREL I.
BIRD P. *see* CALVERT J.E.
BIRD P. *see* DEVEY M.E.
BISCHOFF P. *see* MOOG C.
BLAAS P., *see* YAMAMOTO H.
BLEASDALE-BARR K.M. *see* DEVEY M.E.
BLOOM E.T. *see* UMEHARA H.
BORCA M.V. *see* SRIKUMARAN S.
BOYD R.L. *see* GODFREY D.I.
BOYD R.L. *see* WILSON T.J.
BRACH M. *see* REIDEL D.
BRONDZ B.D. *see* ZAITSEVA M.B.
BROUWER M. *see* DE JONG R.
BUJDOSO R. *see* DUTIA B.M.
BUSSE R. *see* HAUSCHILDT S.
BUTTKE T.M. *see* TEBBEY P.W.

CAHILL R.N.P. *see* WITHERDEN D.A.
CALVERT J.E., JOHNSTONE R., DUGGAN-KEEN M.F. & BIRD P. Immunoglobulin G subclasses secreted by human B cells *in vitro* in response to interleukin-2 and polyclonal activators, 162
CAMPBELL P.A. *see* STEDMAN K.E.
CARLISLE M.S., MCGREGOR D.D. & APPLETON J.A. The role of mucus in antibody-mediated rapid expulsion of *Trichinella spiralis* in suckling rats, 126
CHAO T.-Y., OHNISHI H. & CHU T.M. Indirect inhibition of generation of murine lymphokine-activated killer cell activity in splenocyte cultures by interferon-gamma, 116
CHAPMAN C.J. *see* JORDAN R.K.
CHAUDRY I.H. *see* AYALA A.
CHEERS C., ZHAN Y.F. & EGAN P.J. *In vivo* IL-1 potentiates both specific and non-specific arms of immune response to infection, 411
CHEERS C. *see* EGAN P.
CHILD J.A. *see* JONES R.A.
CHOW S.C. & JONDAL M. A central role for phosphoinositide hydrolysis in activating the lytic mechanisms of human natural killer cells, 106
CHU T.M. *see* CHAO T.-Y.
COMPSTON D.A.S. *see* SCOLDING N.J.
CROATTO M. *see* PHILLIPS W.A.

DAHA M.R. *see* SPECHT GRIJP-GLANDORF R.J.A.M.
DAVÉ A. *see* BATE C.A.W.
DAVRINCHE C. *see* DEMARES M.J.
DAY M.J. *see* MACPHEE I.A.M.
DEACON N.J. *see* PURCELL D.F.J.
DE HEER E. *see* SPECHT GRIJP-GLANDORF R.J.A.M.
DE JONG R., BROUWER M., REBEL V.I., VAN SEVENTER G.A., MIEDEMA F. & VAN LIER R.A.W. Generation of alloreactive cytolytic T lymphocytes by immobilized anti-CD3 monoclonal antibodies. Analysis of requirements for human cytolytic T-lymphocyte differentiation, 357
DEKKER-NOOREN C.C.E.M. *see* SPECHT GRIJP-GLANDORF R.J.A.M.
DEMARES M.J. & DAVRINCHE C. Prevalence of the 1·8 kb complement factor H mRNA in human lung, 150
DEVEY M.E., BLEASDALE-BARR K.M., BIRD P. & AMLOT P.L. Antibodies of different human IgG subclasses show distinct patterns of affinity maturation after immunization with keyhole limpet haemocyanin, 168
DING L. *see* ZHANG Y.H.
DOE W.F. *see* PAVLI P.
DOUGAN G. *see* TITE J.P.
DUGGAN-KEEN M.F. *see* CALVERT J.E.
DUTTA B.M., HOPKINS J., ALLINGTON M.P., BUJDOSO R. & MCCONNELL I. Characterization of monoclonal antibodies specific for α- and β-chains of sheep MHC class II, 27

EARLY E. *see* LAURENCE J.
EGAN P. & CHEERS C. Relationship between colony-stimulating activity and interferon production during infection, 191
EGAN P.J. *see* CHEERS C.
EGIDO J. *see* GONZÁLEZ-CABRERO J.
ELZENGA-CLAASEN I. *see* SLUITER W.

FEHLMANN M. *see* AUBERGER P.
FERNANDEZ-RUIZ E. *see* SOMOZA C.
FERRY B.L., STARKEY P.M., SARGENT I.L., WATT G.M.O., JACKSON M. & REDMAN C.W.G. Cell populations in the human early pregnancy decidua: natural killer activity and response to interleukin-2 of CD56-positive large granular lymphocytes, 446
FINBLOOM D.S. *see* PURI R.K.

FITZPATRICK D.R., SNIDER M., MCDOUGALL L., BESKORWAYNE T., BABIUK L.A., ZAMB T.J. & BEILEFELDT OHMANN H. Molecular mimicry: a herpesvirus glycoprotein antigenically related to a cell-surface glycoprotein expressed by macrophages, polymorphonuclear leucocytes and platelets, 504
FUKUMOTO T. *see* YAMASHITA A.

GALLART T. *see* LOZANO F.
GAO X.-M. *see* TITE J.P.
GENTILE T. *see* MALAN BOREL I.
GODFREY D.I., IZON D.J., TUČEK C.L. WILSON T.J. & BOYD R.L. The phenotypic heterogeneity of mouse thymic stromal cells, 66
GONZALÉZ E. *see* GONZÁLEZ-CABRERO J.
GONZÁLEZ-CABRERO J., EGIDO J., BARAT A. & GONZÁLEZ E. Detection and characterization of circulating and glomerular immune complexes in experimental IgA nephropathy, 296
GONZALEZ-CRUSSI F. *see* HSUEH W.
GOWDA T.V. *see* KASTURI S.
GRAHAM C.M. *see* BARNETT B.C.
GREGG E.O., YARWOOD L., WAGSTAFFE M.J., PEPPER D.S. & MACDONALD M.C. Immunomodulatory properties of platelet factor 4: prevention of concanavalin A suppressor-induction *in vitro* and augmentation of an antigen-specific delayed-type hypersensitivity response *in vivo*, 230
GRIFFITHS M.R. *see* SANDILANDS G.P.
GROSS O. *see* SREDNI B.
GURWITH M.J. *see* SREDNI B.

HALLETT M.B. *see* Petersen M.
HAMILTON J.A. *see* PHILLIPS W.A.
HÄNSCH G.M. *see* YAMAMOTO H.
HARTLMAYR I. *see* BARNETT B.C.
HARVEY J.E. & JONES D.B. Distribution of LCA protein subspecies and the cellular adhesion molecules LFA-1, ICAM-1 and p150,95 within human foetal thymus, 203
HATTORI Y. *see* YAMASHITA A.
HAUSCHILDT S., BASSENGE E., BESSLER W., BUSSE R. & MÜLSCH A. L-arginine-dependent nitric oxide formation and nitrite release in bone marrow-derived macrophages stimulated with bacterial lipopeptide and lipopolysaccharide, 332
HEIN W.R. *see* ALEKSANDERSEN M.
HELLSTRÖM U.B. *see* SYLVAN S.P.E.
HERRMANN F. *see* REIDEL D.
HEYWORTH M.F. & PAPPO J. Recognition of a 30,000 MW antigen of *Giardia muris* trophozoites by intestinal IgA from *Giardia*-infected mice, 535
HOLÁŇ V. *see* LIPOLDOVÁ M.
HONDA H., KIMURA H. & ROSTAMI A. Demonstration and phenotypic characterization of resident macrophages in rat skeletal muscle, 272
HOPKINS J. *see* DUTIA B.M.
HORMAECHE C.E., MASTROENI P., ARENA A., UDDIN J. & JOYSEY H.S. T cells do not mediate the initial suppression of a salmonella infection in the RES, 247
HSUEH W., SUN X., RIOJA L.N. & GONZALEZ-CRUSSI F. The role of the complement system in shock and tissue injury induced by tumour necrosis factor and endotoxin, 309
HUGHES-JENKINS C.M. *see* TITE J.P.
HUME D.A. *see* PAVLI P.

IIKURA Y. *see* MOQBEL R.
INOUE T. *see* YOSHIKAI Y.
ISHII A. *see* KIMURA J.Y.
ISHIZAKA A., JOH K., SHIBATA R., WAGATSUMA Y., NAKANISHI M., TOMIZAWA K., KOJIMA K., KANDIL E., SAKIYAMA Y. & MATSUMOTO S. Regulation of IgE and IgG4 synthesis in patients with hyper IgE syndrome, 414
ISOBE K. *see* ZHANG Y.H.
IZON D.J. *see* GODFREY D.I.

JACKSON M. *see* FERRY B.L.
JAY F.T. *see* LEE C.L.Y.
JI Y.H. *see* MOOG C.
JOH K. *see* ISHIZAKA A.
JOHNSON P.M. *see* PURCELL D.F.J.
JOHNSTONE R. *see* CALVERT J.E.
JONDAL M. *see* CHOW S.C.
JONES D.B. *see* HARVEY J.E.
JONES J. *see* SCOLDING N.J.
JONES R.A., RICHARDS S.J., ROBERTS B.E., CHILD J.A. & SCOTT C.S. Phenotypic switch from CD45RA$^+$ to CD45RA$^-$ by normal blood T cells is associated with increased HLA-ABC expression for CD4$^+$ and CD8$^+$ populations but not for the NK-associated CD4$^-$CD8^{dim+} or CD4$^-$ CD8$^-$ fractions, 55
JORDAN R.K., CHAPMAN C.J., MCLACHLAN S.M., MILNE D., LOVELAND B.E. & ROBINSON J.H. IgG antibody and delayed-type hypersensitivity responses in nude mice grafted with thymic epithelium, 453
JOYSEY H.S. *see* HORMAECHE C.E.
JUSTEMENT L.B. *see* STEDMAN K.E.

KALECHMAN Y. *see* SREDNI B.
KANDIL E. *see* ISHIZAKA A.
KASAHARA Y. *see* YOKOI T.
KASTURI S., RUDRAMMAJI L.M.S. & GOWDA T.V. Antibodies to a phospholipase A_2 from *Vipera russelli* selectively neutralize venom neurotoxicity, 175
KATO K. *see* YOKOI T.
KATOH M. *see* NOSE M.
KAY A.B. *see* MOQBEL R.
KAY N.E., MORLEY J.E. & ALLEN J.I. Interaction between endogenous opioids and IL-2 on PHA-stimulated human lymphocytes, 485
KIMPTON W.G. *see* WITHERDEN D.A.
KIMURA H. *see* HONDA H.
KIMURA J.Y., OHTA N., ISHII A., NAGANO T. & USUI M. Functional characterization of lymphocyte response to fractionated house dust mite antigens (*Dermatophagoides pteronyssinus*) in atopic and non-atopic individuals, 385
KISHIHARA K. *see* OHGA S.
KOJIMA K. *see* ISHIZAKA A.
KÖLLER M. & KÖNIG W. Arachidonic acid metabolism in heat-shock treated human leucocytes, 458
KÖNIG W. *see* KÖLLER M.
KOTANI M. *see* YAMASHITA A.
KRILIS S. *see* ATKINSON Y.H.
KUNTZ L., VELIN D., PFLUMIO F. & LOOR F. Adoptive transfer of viable motheaten humoral autoimmunity in cyclophosphamide-immunodepressed beige recipient mice, 520
KYOGOKU M. *see* NOSE M.

LANDSVERK T. *see* ALEKSANDERSEN M.
LAURENCE J., SAUNDERS A., EARLY E. & SALMON J.E. Human immunodeficiency virus infection of monocytes: relationship to Fc-gamma receptors and antibody-dependent viral enhancement, 338
LEE C.L.Y., LEE S.H.S., JAY F.T. & ROZEE K.R. Immunobiological study of interferon-gamma-producing cells after staphylococcal enterotoxin B stimulation, 94
LEE S.H.S. *see* LEE C.L.Y.
LELAND P. *see* PURI R.K.
LIEW F.Y. *see* TITE J.P.
LINDEMANN A. *see* REIDEL D.
LIPOLDOVÁ M. & HOLÁŇ V. Expression of genes for interleukin-1α and tumour necrosis factor-α in newborn mice, 136
LIPSCOMBE M. *see* TITE J.P.
LOOR F. *see* KUNTZ L.
LOPEZ A.F. *see* ATKINSON Y.H.
LOVELAND B.E. *see* JORDAN R.K.
LOZANO F., ALBEROLA-ILA J., PLACES L., GALLART T. & VIVES J. Protein kinase C-dependent up-regulation of CD5 surface expression on normal and lymphoblastoid T cells, 434
LUBLIN D.M. *see* PURCELL D.F.J.
LUU B. *see* MOOG C.

MCCLURE S. *see* ALEKSANDERSEN M.
MCCONNELL I. *see* DUTIA B.M.
MACDONALD A.J. *see* MOQBEL R.
MACDONALD M.C. *see* GREGG E.O.
MCDOUGALL L. *see* FITZPATRICK D.R.
MCGREGOR D.D. *see* CARLISLE M.S.
MACHUGH N.D. *see* TOYE P.G.

McKenzie I.F.C. *see* Purcell D.F.J.
McLachlan S.M. *see* Jordan R.K.
MacPhee I.A.M., Day M.J. & Mason D.W. The role of serum factors in the suppression of experimental allergic encephalomyelitis: evidence for immunoregulation by antibody to the encephalitogenic peptide, 527
MacPhee I. *see* Mason D.
Maghazachi A.A. Differential effects of various cytokines on the generation of rat LAK cells from their purified precursors, 465
Malan Borel I., Gentile T., Angelucci J., Margni R.A. & Binaghi R.A. Asymmetric Fab glycosylation in guinea-pig IgG1 and IgG2, 281
Margni R.A. *see* Malan Borel I.
Mari B. *see* Auberger P.
Mason D., MacPhee I. & Antoni F. The role of the neuroendocrine system in determining genetic susceptibility to experimental allergic encephalomyelitis in the rat, 1
Mason D. & Powrie F. Memory $CD4^+$ T cells in man form two distinct subpopulations, defined by their expression of isoforms of the leucocyte common antigen, CD45, 427
Mason D.W. *see* MacPhee I.A.M.
Mastroeni P. *see* Hormaeche C.E.
Matsumoto S. *see* Ishizaka A.
Matsuzaki G. *see* Ohga S.
Matsuzaki G. *see* Yoshikai Y.
Matthews N. *see* Petersen M.
Matthews N. *see* Steadman R.
Mayrhofer G. & Spargo L.D.J. Distribution of class II major histocompatibility antigens in enterocytes of the rat jejunum and their association with organelles of the endocytic pathway, 11
Mertelsmann R. *see* Reidel D.
Michlin H. *see* Sredni B.
Miedema F. *see* De Jong R.
Milne D. *see* Jordan R.K.
Miyasaka M. *see* Yamashita A.
Miyauchi Y. *see* Shimokawa Y.
Miyawaki T. *see* Yokoi T.
Moog C., Ji Y.H., Waltzinger C., Luu B. & Bischoff P. Studies on the immunological properties of oxysterols: *in vivo* actions of 7,25-dihydroxycholesterol upon murine peritoneal cells, 344
Moqbel R., Walsh G.M., Nagakura T., MacDonald A.J., Wardlaw A.J., Iikura Y. & Kay A.B. The effect of platelet-activating factor on IgE binding to, and IgE-dependent biological properties of, human eosinophils, 251
Morgan B.P. *see* Scolding N.J.
Morley J.E. *see* Kay N.E.
Morrison W.I. *see* Toye P.
Mostowski H. *see* Puri R.K.
Mülsch A. *see* Hauschildt S.
Murray A.W. *see* Atkinson Y.H.
Muthukkaruppan V.R. *see* Vijayakumar R.K.

Nagakura T. *see* Moqbel R.
Nagano T. *see* Kimura J.Y.
Nagase F. *see* Zhang Y.H.
Nakanishi M. *see* Ishizaka A.
Nakano K. *see* Okamoto H.
Nakashima I. *see* Zhang Y.H.
Nataraj C. *see* Srikumaran S.
Nibbering P.H. *see* Sluiter W.
Nicholson-Weller A. *see* Yamamoto H.
Nomoto K. *see* Ohga S.
Nomoto K. *see* Yoshikai Y.
Nose M., Katoh M., Okada N., Kyogoku M. & Okada H. Tissue distribution of HRF20, a novel factor preventing the membrane attack of homologous complement, and its predominant expression on endothelial cells *in vivo*, 145

O'Callaghan D. *see* Tite J.P.
Ogimoto M. *see* Ohga S.
Oglesby T.J. *see* Purcell D.F.J.
Oh C. *see* Okamoto H.
Ohga S., Yoshikai Y., Kishihara K., Matsuzaki G., Ogimoto M. & Nomoto K. Different expression of T-cell receptor β-chain variable region genes in lymph nodes of lpr mice with different alleles of the major histocompatibility complex, 216
Ohnishi H. *see* Chao T.-Y.
Ohno I., Tanno Y., Yamauchi K. & Takishima T. Gene expression and production of tumour necrosis factor by a rat basophilic leukaemia cell line (RBL-2H3) with IgE receptor triggering, 88
Ohta N. *see* Kimura J.Y.
Okada H. *see* Nose M.
Okada N. *see* Nose M.
Okamoto H., Oh C. & Nakano K. Possible involvement of adenosine 3′:5′-cyclic monophosphate and extracellular calcium ions in histamine stimulation of interleukin-1 release from macrophage-like P388D1 cells, 186
Onisk D.V. *see* Srikumaran S.

Pappo J. *see* Heyworth M.F.
Parr E.L. *see* Thapar M.A.
Parr M.B. *see* Thapar M.A.
Pavli P., Woodhams C.E., Doe W.F. & Hume D.A. Isolation and characterization of antigen-presenting dendritic cells from the mouse intestinal lamina propria, 40
Pemberton R.M., Wraith D.C. & Askonas B.A. Influenza peptide-induced self-lysis and down-regulation of cloned cytotoxic T cells, 223
Pepper D.S. *see* Gregg E.O.
Perrin M.M. *see* Ayala A.
Petersen M., Steadman R., Hallett M.B., Matthews N. & Williams J.D. Zymosan-induced leukotriene B_4 generation by human neutrophils is augmented by rhTNF-α but not chemotactic peptide, 75
Petersen M.M. *see* Steadman R.
Petrycky L.D. *see* Rayner D.C.
Peyron J.-F. *see* Auberger P.
Pflumio F. *see* Kuntz L.
Phillips W.A., Croatto M. & Hamilton J.A. Priming the macrophage respiratory burst with IL-4: enhancement with TNF-α but inhibition by IFN-γ, 498
Places L. *see* Lozano F.
Playfair J.H.L. *see* Bate C.A.W.
Powrie F. *see* Mason D.
Pugh-Humpreys R.G.P., Ross C.S.K. & Thomson A.W. The influence of FK-506 on the thymus: an immunophenotypic and structural analysis, 398
Purcell D.F.J., McKenzie I.F.C., Lublin D.M., Johnson P.M., Atkinson J.P., Oglesby T.J. & Deacon N.J. The human cell-surface glycoproteins HuLy-m5 membrane co-factor protein (MCP) of the complement system, and trophoblast leucocyte-common (TLX) antigen, are CD46, 155
Puri R.K., Finbloom D.S., Leland P., Mostowski H. & Siegel J.P. Expression of high-affinity IL-4 receptors on murine tumour infiltrating lymphocytes and their up-regulation by IL-2, 492

Rahman S.M.J. *see* Zhang Y.H.
Ramírez F. *see* Somoza C.
Rayner D.C. & Petrycky L.D. Class II-restricted bifunctional T-cell hybridomas reactive to self- and foreign myelin basic protein, 365
Rebel V.I. *see* De Jong R.
Rebollo A. *see* Somoza C.
Redman C.W.G. *see* Ferry B.L.
Reidel D., Lindemann A., Brach M., Mertelsmann R. & Herrmann F. Granulocyte–macrophage colony-stimulating factor and interleukin-3 induce surface expression of interleukin-2 receptor p55-chain and CD4 by human eosinophils, 258
Richards S.J. *see* Jones R.A.
Richards S.J. *see* Scott C.S.
Rioja L.N. *see* Hsueh W.
Roberts B.E. *see* Jones R.A.
Roberts B.E. *see* Scott C.S.
Robinson J.H. *see* Jordan R.K.
Ross C.S.K. *see* Pugh-Humpreys R.G.P.
Rostami A. *see* Honda H.
Rozee K.R. *see* Lee C.L.Y.
Rudrammaji L.M.S. *see* Kasturi S.

Sakiyama Y. *see* Ishizaka A.
Salmon J.E. *see* Laurence J.

SANDILANDS G.P., AHMED A.E.E., GRIFFITHS M.R. & WHALEY K. Immunohistochemical localization of a plasma protein (glycoprotein 60) which inhibits complement-mediated prevention of immune precipitation, 303
SANZ E. *see* SOMOZA C.
SARGENT I.L. *see* FERRY B.L.
SAUNDERS A. *see* LAURENCE J.
SAVIC V., STEFANOVIC V., ARDAILLOU N. & ARDAILLOU R. Induction of ecto-5′-nucleotidase of rat cultured mesangial cells by interleukin-1β and tumour necrosis factor-α, 321
SCOLDING N.J., JONES J., COMPSTON D.A.S. & MORGAN B.P. Oligodendrocyte susceptibility to injury by T-cell perforin, 6
SCOTT C.S., RICHARDS S.J. & ROBERTS B.E. Patterns of membrane TcR$\alpha\beta$ and TcR$\gamma\delta$ chain expression by normal blood $CD4^+CD8^-$, $CD4^-CD8^+$, $CD4^-CD8^{dim+}$ and $CD4^-CD8^-$ lymphocytes, 351
SCOTT C.S. *see* JONES R.A.
SEHGAL S.N. *see* SREDNI B.
SHARON P. *see* SREDNI B.
SHIBATA R. *see* ISHIZAKA A.
SHIMOKAWA Y., TAKEYA M., MIYAUCHI Y. & TAKAHASHI K. A monoclonal antibody, RbM2, specific for a lysosomal membrane antigen of rabbit monocyte/macrophages, 513
SIEGEL J.P. *see* PURI R.K.
SILVA A. *see* SOMOZA C.
SLUITER W., NIBBERING P.H., VAN FURTH R., ANNEMA A., ELZENGA-CLAASEN I., VAN HEMSBERGEN-OOMENS L.W.M. & VAN DER VOORT VAN DER KLEIJ-VAN ANDEL. Increased activity of FIM in serum of mice during a *Mycobacterium bovis* (BCG) infection, 327
SNIDER M. *see* FITZPATRICK D.R.
SOMOZA C., FERNÁNDEZ-RUIZ E., REBOLLO A., SANZ E., RAMÍREZ F. & SILVA A. OX48, a monoclonal antibody against a 70,000 MW activation antigen expressed by T cells bearing the high-affinity interleukin-2 receptor, 210
SONTHONNAX S. *see* AUBERGER P.
SPARGO L.D.J. *see* MAYRHOFER G.
SPECHT GRIJP-GLANDORF R.J.A.M., DE HEER E., DEKKER-NOOREN C.C.E.M., DAHA M.R. & VAN ES L.A. Specificity and cross-reactive idiotypes of anti-gp330 autoantibodies in active Heymann nephritis, 290
SREDNI B., KALECHMAN Y., ALBECK M., GROSS O., AURBACH D., SHARON P., SEHGAL S.N., GURWITH M.J. & MICHLIN H. Cytokine secretion effected by synergism of the immunomodulator AS101 and the protein kinase C inducer bryostatin, 473
SRIKUMARAN S., ONISK D.V., BORCA M.V., NATARAJ C. & ZAMB T.J. Anti-idiotypic antibodies induce neutralizing antibodies to bovine herpesvirus 1, 284
STARKEY P.M. *see* FERRY B.L.
STEADMAN R., PETERSEN M.M., WILLIAMS D., MATTHEWS N. & WILLIAMS J.D. The selective augmentation by recombinant human tumour necrosis factor-alpha of neutrophil responses to pathogenic *Escherichia coli*, 133
STEADMAN R. *see* PETERSEN M.
STEDMAN K.E., JUSTEMENT L.B. & CAMPBELL P.A. Endogenously secreted IL-4 is required for mouse thymocytes to become cytotoxic. Human, but not mouse, IL-2 induces a functionally immature thymic subset to secrete IL-4 and become CTL, 478
STEFANOVIC V. *see* SAVIC V.
SUN X. *see* HSUEH W.
SYLVAN S.P.E. & HELLSTRÖM U.B. HBsAg-induced interferon-gamma secretion in T cells form asymptomatic HBsAg carriers and HB-immune donors *in vitro*, 197

TAKAHASHI K. *see* SHIMOKAWA Y.
TAKEYA M. *see* SHIMOKAWA Y.
TAKISHIMA T. *see* OHNO I.
TANIGUCHI N. *see* YOKOI T.
TANNO Y. *see* OHNO I.
TAVERNE J. *see* BATE C.A.W.
TEALE A.J. *see* TOYE P.G.
TEBBEY P.W. & BUTTKE T.M. Molecular basis for the immunosuppressive action of stearic acid on T cells, 379
THAPAR M.A., PARR E.L. & PARR M.B. Secretory immune responses in mouse vaginal fluid after pelvic, parenteral or vaginal immunization, 121
THOMAS D.B. *see* BARNETT B.C.
THOMSON A.W. *see* PUGH-HUMPREYS R.G.P.
TITE J.P. Differential requirement for protein synthesis in cytolysis mediated by class I and class II MHC-restricted cytotoxic T cells, 440
TITE J.P., GAO X.-M., HUGHES-JENKINS C.M., LIPSCOMBE M., O'CALLAGHAN D., DOUGAN G. & LIEW F.Y. Anti-viral immunity induced by recombinant nucleoprotein of Influenza A virus. III. Delivery of recombinant nucleoprotein to the immune system using attenuated *Salmonella typhimurium* as a live carrier, 540
TOMIZAWA K. *see* ISHIZAKA A.
TOYE P.G., MACHUGH N.D., BENSAID A.M., ALBERTI S., TEALE A.J. & MORRISON W.I. Transfection into mouse L cells of genes encoding two serologically and functionally distinct bovine class I MHC molecules from a MHC-homozygous animal: evidence for a second class I locus in cattle, 20
TUČEK C.L. *see* GODFREY D.I.

UDDIN J. *see* HORMAECHE C.E.
UMEHARA H. & BLOOM E.T. The IL-2 receptor beta subunit is absolutely required for mediating the IL-2-induced activation of NK activity and proliferative activity of human large granular lymphocytes, 111
USUI M. *see* KIMURA J.Y.

VADAS M.A. *see* ATKINSON Y.H.
VAN DER VOORT VAN DER KLEIJ-VAN ANDEL *see* SLUITER W.
VAN ES L.A. *see* SPECHT GRIJP-GLANDORF R.J.A.M.
VAN FURTH R. *see* SLUITER W.
VAN HEMSBERGEN-OOMENS L.W.M. *see* SLUITER W.
VAN LIER R.A.W. *see* DE JONG R.
VAN SEVENTER G.A. *see* DE JONG R.
VEITCH J. *see* AHMED A.E.E.
VELIN D. *see* KUNTZ L.
VIJAYAKUMAR R.K. & MUTHUKKARUPPAN V.R. Analysis of carrageenan-induced suppression of antibody response, 181
VIVES J. *see* LOZANO F.

WAGATSUMA Y. *see* ISHIZAKA A.
WAGSTAFFE M.J. *see* GREGG E.O.
WALSH G.M. *see* MOQBEL R.
WALTZINGER C. *see* MOOG C.
WARDLAW A.J. *see* MOQBEL R.
WASHINGTON E.A. *see* WITHERDEN D.A.
WATT G.M.O. *see* FERRY B.L.
WHALEY K. *see* AHMED A.E.E.
WHALEY K. *see* SANDILANDS G.P.
WILLIAMS D. *see* STEADMAN R.
WILLIAMS J.D. *see* PETERSEN M.
WILLIAMS J.D. *see* STEADMAN R.
WILSON T.J. & BOYD R.L. Cyclophosphamide- and testosterone-induced alteration in chicken bursal stroma identified by monoclonal antibodies, 241
WILSON T.J. *see* GODFREY D.I.
WITHERDEN D.A., KIMPTON W.G., WASHINGTON E.A. & CAHILL R.N.P. Non-random migration of $CD4^+$, $CD8^+$ and $\gamma\delta^+T19^+$ lymphocytes through peripheral lymph nodes, 235
WOODHAMS C.E. *see* Pavli P.
WRAITH D.C. *see* PEMBERTON R.M.

YACHIE A. *see* YOKOI T.
YAMAMOTO H., BLAAS P., NICHOLSON-WELLER A. & HANSCH G.M. Homologous species restriction of the complement-mediated killing of nucleated cells, 422
YAMASHITA A., HATTORI Y., KOTANI M., MIYASAKA M. & FUKUMOTO T. A novel monoclonal antibody, Mar 1, directed specifically against mononuclear phagocyte system cells in rats, 262
YAMAUCHI K. *see* OHNO I.
YARWOOD L. *see* GREGG E.O.
YOKOI T., MIYAWAKI T., YACHIE A., KATO K., KASAHARA Y. & TANIGUCHI N. Epstein–Barr virus-immortalized B cells produce IL-6 as an autocrine growth factor, 100
YOSHIDA T. *see* ZHANG Y.H.
YOSHIKAI Y., MATSUZAKI G., INOUE T. & NOMOTO K. An increase in number of T-cell receptor γ/δ-bearing T cells in athymic nude mice treated with complete Freund's adjuvants, 61
YOSHIKAI Y. *see* OHGA S.

ZAITSEVA M.B. & BRONDZ B.D. Mechanism of MHC class II restriction in the interaction between specific suppressor and responder T cells in a proliferative response: Ia interaction with a putative anti-self receptor, expressed on pre-activated responder cells, 372
ZAMB T.J. *see* FITZPATRICK D.R.
ZAMB T.J. *see* SRIKUMARAN S.
ZHAN Y.F. *see* CHEERS C.
ZHANG Y.H., YOSHIDA T., ISOBE K., RAHMAN S.M.J., NAGASE F., DING L. & NAKASHIMA I. Modulation by glycyrrhizin of the cell-surface expression of H-2 class I antigens on murine tumour cell lines and normal cell populations, 405

Subject Index

Activation antigen, rat:
  expressed by IL-2R T cells, 210
Affinity maturation:
  different IgG subclasses, 168
Alloreactive CTL:
  generated by anti-CD3, 357
Antibody-mediated expulsion of *T. spiralis*:
  role of mucus, 126
Anti-idiotype antibodies:
  induce neutralizing antibodies to bovine herpesvirus 1, 284
Anti-neurotoxic antibodies to *V. russelli* venom, 175
Arachidonic acid metabolism:
  in heat-shock treated leucocytes, 458
Autoimmunity, humoral:
  in me^v/bg B6 chimaeric mice, 520

Bursal stroma, chicken:
  alterations induced by cyclophosphamide and testosterone, 241

Carrageenan:
  suppression of antibody response, 181
$CD4^+$ T cells, human:
  subpopulations defined by expression of CD45, 427
CD5 expression:
  protein kinase C-dependent up-regulation, 434
CD46:
  includes HuLy-m5 MCP and TLX antigens, 155
Class II antigen:
  in rat jejunal epithelium, 11
Colony stimulation and IFN production during infection, 191
Complement factor H:
  1·8 kb mRNA in human lung, 150
Complement-mediated immune precipitation:
  inhibition, 139
Complement-mediated killing:
  nucleated cells, species restriction, 422
Complement-mediated prevention of immune precipitation:
  inhibition by glycoprotein 60, 303
Cytokine secretion:
  effect of synergism of AS101 and bryostatin, 473
Cytolysis, mediated by class I- and II-restricted T cells:
  differential requirement for protein synthesis, 440
Cytotoxic T cells:
  influenza peptide-induced self-lysis and down-regulation, 223
Cytotoxic thymocytes:
  development and IL-4, 478

Dendritic cells, antigen-presenting:
  from intestinal lamina propria, 40
Dermatophagoides pteronyssinus:
  characterization of lymphocyte response to fractionated antigens in atopic and non-atopic individuals, 385

Ecto-5′-nucleotidase:
  induced by IL-1β and TNF-α, 321
Eosinophils:
  effect of platelet-activating factor, 251
Eosinophils, human:
  expression of IL-2R p55 and CD4, 258
Experimental allergic encephalomyelitis:
  role of neuroendocrine system, 1
  suppression, role of serum factors, 527

FIM, factor increasing monocytopoiesis:
  and BCG infection, 327
FK-506:
  effects on the thymus, 398

Giardia antigen:
  recognized by intestinal IgA, 535
Glycoprotein 60:
  histochemical localization, 303
Glycosylation, asymmetric:
  in guinea-pig Ig, 281

H-2 class I antigen expression:
  modulation by glycyrrhizin, 405
Heat-shock treated leucocytes:
  arachidonic acid metabolism, 458
HLA-ABC expression:
  2H4 defined, 55
HRF20:
  tissue distribution, 145
Human immunodeficiency virus:
  Fc-gamma receptors and antibody-dependent enhancement, 338
Hyper IgE syndrome:
  regulation of IgE and IgG4 synthesis, 414

Idiotypes, cross-reactive:
  in Heymann nephritis, 290
Immune complexes:
  in IgA nephropathy, 296
Immunoglobulin A nephropathy:
  detection and characterization of immune complexes, 296
Immunoglobulin G subclasses:
  affinity maturation, 168
Immunoglobulin G subclasses, human:
  in vitro, 162
Immunosuppression:
  by stearic acid, 379
Influenza haemagglutinin:
  $I\text{-}A^d$- and $I\text{-}E^d$-restricted T-cell epitopes, 48
Interferon production and colony-stimulating activity during infection, 191
Interferon-gamma-producing cells:
  after enterotoxin B stimulation, 94
Interferon-gamma production in asymptomatic HBsAg carriers, 197
Interleukin-1:
  potentiation of non-specific and specific immune response, 411
Interleukin-1 and tumour necrosis factor:
  expression of genes in newborn mice, 136
Interleukin-1 release:
  involvement of cyclic AMP and extracellular calcium, 186
Interleukin-2 production:
  requirement for chymotryptic-type serine protease, 547
Interleukin-2 receptor beta subunit:
  in IL-2-regulated NK cells, 111
Interleukin-4:
  priming of macrophage respiratory burst, 498
  required for development of cytotoxic thymocytes, 478
Interleukin-4 receptors, high affinity: up-regulation by IL-2, 492
Interleukin-6:
  produced by EBV-immortalized B cells, 100

Large granular lymphocytes:
  in early pregnancy decidua, 446

Leukotriene B_4:
generation by human neutrophils, 75
Lymphocyte subsets:
distribution in large intestinal follicles of lambs, 391
Lymphokine-activated killer cells:
generation inhibited by IFN-γ, 116
Lymphokine-activated killer cells, rat:
effects of cytokines on development, 465
Lysosmal membrane antigen:
rabbit monocyte/macrophage, monoclonal antibody to, 513

Macrophage antigen presentation:
after haemorrhage, 33
Macrophages, in rat skeletal muscle:
characterization, 272
Major histocompatibility complex class II restriction:
in interactions between suppressor and responder T cells, 372
Major histocompatibility complex class II, sheep:
characterization of monoclonal antibodies specific for α- and β-chains, 27
Malaria exoantigens:
induction of antibody which blocks TNF production, 315
Migration, non-random:
of T-lymphocyte subsets, 235
Molecular mimicry:
herpesvirus glycoprotein, 504
Mononuclear phagocyte stem cells:
monoclonal antibody directed to, 262
Mucus:
role of antibody-mediated expulsion of *T. spiralis*, 126

Natural killer cells, lytic mechanisms:
role of phosphoinositide hydrolysis, 106
Nitric oxide and nitrite:
in bone marrow-derived macrophages, 332
Nucleoprotein:
delivery by a live *S. typhimurium* carrier, 540

Opioids and IL-2:
interaction on PHA-stimulated human lymphocytes, 485
Oxysterols:
in vivo action on peritoneal cells, 344

Perforin, T cell:
oligodendrocyte susceptibility to, 6
Platelet-activating factor:
effect on IgE binding to eosinophils, 251
Platelet factor 4:
immunomodulatory properties, 230

Secretory immune responses in mouse vaginal fluid, 121

T-cell hybridomas, MBP-specific:
class II restricted, 365
T-cell receptor chain expression, 351
T-cell receptor V genes: in lpr mice, 216
T cells:
in complete Freund's adjuvant-treated nude mice, 61
role in suppression of salmonella infection, 247
Thymic epithelial grafts:
in nude mice, 453
Thymic stromal cells:
phenotypic heterogeneity, 66
Thymus, foetal:
distribution of LCA and adhesion molecules, 203
Transfection of bovine class I MHC, 20
Tumour necrosis factor:
augmentation of neutrophil responses, 133
gene expression and production, 88
shock and tissue injury, role of complement, 309
Tumour necrosis factor, human:
stimulates arachidonic acid release from neutrophils, 82